

POWERS OF DISTANCES TO LOWER DIMENSIONAL SETS AS MUCKENHOUP WEIGHTS

H. AIMAR¹, M. CARENA^{1,2,*}, R. DURÁN³ and M. TOSCHI^{1,†}

¹Instituto de Matemática Aplicada del Litoral (CONICET-UNL),
Departamento de Matemática (FIQ-UNL), Santa Fe, Argentina
e-mails: haimar@santafe-conicet.gov.ar, mtoschi@santafe-conicet.gov.ar

²Instituto de Matemática Aplicada del Litoral (CONICET-UNL),
Departamento de Matemática (FHUC-UNL), Santa Fe, Argentina
e-mail: mcarena@santafe-conicet.gov.ar

³Instituto de Investigaciones Matemáticas "Luis A. Santaló" (CONICET-UBA),
Departamento de Matemática (UBA), Buenos Aires, Argentina
e-mail: rduran@dm.uba.ar

(Received June 14, 2013; revised September 3, 2013; accepted September 3, 2013)

Abstract. Let (X, d, μ) be an Ahlfors metric measure space. We give sufficient conditions on a closed set $F \subseteq X$ and on a real number β in such a way that $d(x, F)^\beta$ becomes a Muckenhoupt weight. We give also some illustrations to regularity of solutions of partial differential equations and regarding some classical fractals.